Author Index

Akins, D.L. and Kumar, V.T.
High-performance liquid chromatography of cyanine dyes. Multiphase separation, purification, and substitution of the counter ion 689(1995)269

Almudaris, A., Ashton, D.S., Ray, A. and Valko, K.
Trace analysis of impurities in 3′-azido-3′-deoxythymidine by reversed-phase high-performance liquid chromatography and thermospray mass spectrometry 689(1995)31

Angulo-Tatis, D., see Oscarsson, S. 689(1995)3

Aoki, F., see Ôi, N. 689(1995)195

Armstrong, D.W., Gasper, M.P. and Rundlett, K.L.
Highly enantioselective capillary electrophoretic separations with dilute solutions of the macrocyclic antibiotic ristocetin A 689(1995)285

Ashton, D.S., see Almudaris, A. 689(1995)31

Aue, W.A., see Singh, H. 689(1995)45

Bächmann, K., see Röder, A. 689(1995)305

Beyermann, M., see Rothemund, S. 689(1995)219

Bienert, M., see Rothemund, S. 689(1995)219

Brenna, J.T., see Goodman, K.J. 689(1995)63

Chaga, G., see Oscarsson, S. 689(1995)3

Chalányová, M., see Hutta, M. 689(1995)123

Chibante, L.P.F., see Heymann, D. 689(1995)157

Conti, M., see Simò-Alfonso, E. 689(1995)85

Dathe, M., see Rothemund, S. 689(1995)219

Desbène, P.L., Morin, C.J., Desbène Monvernay, A.M. and Groult, R.S.
Utilization of fluorescein sodium salt in laser-induced indirect fluorimetric detection of ions separated by capillary zone electrophoresis 689(1995)135

Desbène, P.L. and Rony, C.M.
Determination by high-performance capillary electrophoresis of alkylaromatics used as bases of sulfonation in the preparation of industrial surfactants 689(1995)107

Desbène Monvernay, A.M., see Desbène, P.L. 689(1995)135

Dona, A.-M. and Verchère, J.-F.
High-performance liquid chromatography of alditols with indirect photometric detection 689(1995)13

Elsener, M., see Koebel, M. 689(1995)164

Engelhardt, H., see Rothemund, S. 689(1995)219

Furusaki, S., see Tsuneda, S. 689(1995)211

Gasper, M.P., see Armstrong, D.W. 689(1995)285

Gelfi, C., Orsi, A., Leoncini, F. and Righetti, P.G.
Fluidified polyacrylamides as molecular sieves in capillary zone electrophoresis of DNA fragments 689(1995)97

Gelfi, C., see Simò-Alfonso, E. 689(1995)85

Glatz, C.E., see Heng, M.H. 689(1995)227

Goodman, K.J. and Brenna, J.T.
High-precision gas chromatography–combustion isotope ratio mass spectrometry at low signal levels 689(1995)63

Goto, J., see Iida, T. 689(1995)77

Groult, R.S., see Desbène, P.L. 689(1995)135

Haken, J.K.
Introduction to Open-Tubular Column Gas Chromatography (by J.V. Hinshaw and L.S. Ettre) (Book Review) 689(1995)313

Heng, M.H. and Glatz, C.E.
Charged fusions for β-galactosidase retention in anion-exchange chromatography 689(1995)227

Herdewijn, P., see Thoithi, G. 689(1995)247

Heymann, D., Chibante, L.P.F. and Smalley, R.E.
Determination of C_{60} and C_{70} fullerenes in geologic materials by high-performance liquid chromatography 689(1995)157

Hirokawa, T., Xia, W. and Kiso, Y.
Isotachophoretic separation of rare earth ions. I. Separation behaviour of yttrium and fourteen lanthanide ions forming complexes with tartaric acid and α-hydroxyisobutyric acid 689(1995)149

Hoogmartens, J., see Thoithi, G. 689(1995)247

Hutta, M., Kaniansky, D., Kovalčíková, E., Marák, J., Chalányová, M., Madajová, V. and Šimuničová, E.
Preparative capillary isotachophoresis as a sample pretreatment technique for complex ionic matrices in high-performance liquid chromatography 689(1995)123

Iida, T., Tazawa, S., Tamaru, T., Goto, J. and Nambara, T.
Gas chromatographic separation of bile acid 3-glucosides and 3-glucuronides without prior deconjugation on a stainless-steel capillary column 689(1995)77

Itoh, H., see Nimura, N. 689(1995)203

Jandera, P. and Urbánek, J.
Comparison of chromatographic behaviour of oligoethylene glycol nonylphenyl ether non-ionic and anionic surfactants in reversed-phase high-performance liquid chromatography 689(1995)255

Kaniansky, D., see Hutta, M. 689(1995)123

Khaskhely, G.Q., see Khuhawar, M.Y. 689(1995)39

Khuhawar, M.Y., Lanjwani, S.N. and Khaskhely, G.Q.
High-performance liquid chromatographic determination of vanadium in crude petroleum oils using bis(salicylaldehyde)tetramethylethylenediimine 689(1995)39

Kim, Y., Park, S., Park, J. and Lee, W.
Detection of benzthiazide by high-performance liquid chromatography–thermospray mass spectrometry 689(1995)170

Kinoshita, T., see Nimura, N. 689(1995)203

Kiso, Y., see Hirokawa, T. 689(1995)149

Kisu, N., see Ôi, N. 689(1995)195

Kitahara, H., see Ôi, N. 689(1995)195

Klick, S.
Evaluation of different injection techniques in the gas chromatographic determination of thermolabile trace impurities in a drug substance 689(1995)69

Koebel, M. and Elsener, M.
Determination of urea and its thermal decomposition products by high-performance liquid chromatography 689(1995)164
Körner, A. and Peter, A.
Alternative methods for the determination of trace amounts of 4-aminomorpholine in molsidomine and linsidomine 689(1995)235
Kovalčíková, E., see Hutta, M. 689(1995)123
Kraak, J.C., see Swart, R. 689(1995)177
Krause, E., see Rothemund, S. 689(1995)219
Kumar, V.T., see Akins, D.L. 689(1995)269
Lanjwani, S.N., see Khuhawar, M.Y. 689(1995)39
Lee, W., see Kim, Y. 689(1995)170
Leoncini, F., see Gelfi, C. 689(1995)97
Liu, Y.-M., see Toyo'oka, T. 689(1995)23
López-Pérez, M.J.
Aqueous Two-Phase Systems (Methods in Enzymology, Vol. 228) (edited by H. Walter and G. Johannson) (Book Review) 689(1995)312
Madajová, V., see Hutta, M. 689(1995)123
Marák, J., see Hutta, M. 689(1995)123
Millier, B., see Singh, H. 689(1995)45
Morin, C.J., see Desbène, P.L. 689(1995)135
Nambara, T., see Iida, T. 689(1995)77
Nimura, N., Itoh, H. and Kinoshita, T.
Diol-bonded silica gel as a restricted access packing forming a binary-layered phase for direct injection of serum for the determination of drugs 689(1995)203
Ôi, N., Kitahara, H., Aoki, F. and Kisu, N.
Direct separation of carboxylic acid enantiomers by high-performance liquid chromatography with amide and urea derivatives bonded to silica gel as chiral stationary phases 689(1995)195
Orsi, A., see Gelfi, C. 689(1995)97
Oscarsson, S., Angulo-Tatis, D., Chaga, G. and Porath, J.
Amphiphilic agarose-based adsorbents for chromatography. Comparative study of adsorption capacities and desorption efficiencies 689(1995)3
Park, J., see Kim, Y. 689(1995)170
Park, S., see Kim, Y. 689(1995)170
Peter, A., see Körner, A. 689(1995)235
Phillips, J.B., see Zhang, M. 689(1995)275
Poppe, H., see Swart, R. 689(1995)177
Porath, J., see Oscarsson, S. 689(1995)3
Ray, A., see Almudaris, A. 689(1995)31
Righetti, P.G., see Gelfi, C. 689(1995)97
Righetti, P.G., see Simò-Alfonso, E. 689(1995)85
Röder, A. and Bächmann, K.
Simultaneous determination of organic and inorganic anions in the sub-μmol/l range in rain water by capillary zone electrophoresis 689(1995)305
Roets, E., see Thoithi, G. 689(1995)247
Rony, C.M., see Desbène, P.L. 689(1995)107
Rothemund, S., Krause, E., Beyermann, M., Dathe, M., Engelhardt, H. and Bienert, M.
Recognition of α-helical peptide structures using high-performance liquid chromatographic retention data for D-amino acid analogues: influence of peptide amphipathicity and of stationary phase hydrophobicity 689(1995)219
Rundlett, K.L., see Armstrong, D.W. 689(1995)285
Saito, K., see Tsuneda, S. 689(1995)211
Schwarzenbach, R.
Perfumery—Practice and Principles (by R.R. Calkin and J.S. Jellinek) (Book Review) 689(1995)175
Simò-Alfonso, E., Conti, M., Gelfi, C. and Righetti, P.G.
Sodium dodecyl sulfate capillary electrophoresis of proteins in entangled solutions of poly(vinyl alcohol) 689(1995)85
Šimuničová, E., see Hutta, M. 689(1995)123
Singh, H., Millier, B. and Aue, W.A.
Polarization, relaxation and unrestrictedly linear response in a bipolar, constant-frequency electron-capture detector 689(1995)45
Smalley, R.E., see Heymann, D. 689(1995)157
Sugo, T., see Tsuneda, S. 689(1995)211
Swart, R., Kraak, J.C. and Poppe, H.
Performance of an ethoxyethylacrylate stationary phase for open-tubular liquid chromatography 689(1995)177
Tamaru, T., see Iida, T. 689(1995)77
Tazawa, S., see Iida, T. 689(1995)77
Thoithi, G., Van Schepdael, A., Herdewijn, P., Roets, E. and Hoogmartens, J.
Liquid chromatographic separation of diamino analogues of 2′- or 3′-deoxyadenosine from adenine on a poly(styrene–divinylbenzene) polymer column 689(1995)247
Toyo'oka, T. and Liu, Y.-M.
High-performance liquid chromatographic resolution of amino acid enantiomers derivatized with fluorescent chiral Edman reagents 689(1995)23
Tsuneda, S., Saito, K., Furusaki, S. and Sugo, T.
High-throughput processing of proteins using a porous and tentacle anion-exchange membrane 689(1995)211
Urbánek, J., see Jandera, P. 689(1995)255
Valko, K., see Almudaris, A. 689(1995)31
Van Schepdael, A., see Thoithi, G. 689(1995)247
Verchère, J.-F., see Dona, A.-M. 689(1995)13
Welch, C.J.
Imprintable brush-type chiral stationary phase 689(1995)189
Xia, W., see Hirokawa, T. 689(1995)149
Zhang, M. and Phillips, J.B.
Applications of multiplex gas chromatography to the determination of organics in solid samples 689(1995)275